

# PROPERTIES OF CERTAIN FAMILIES OF $2k$ -CYCLE FREE GRAPHS<sup>1</sup>

F. LAZEBNIK

*Department of Mathematical Sciences  
University of Delaware, Newark, DE 19716, USA*

V. A. USTIMENKO

*Department of Mathematics and Mechanics  
University of Kiev, Kiev 252127, Ukraine*

A. J. WOLDAR

*Department of Mathematical Sciences  
Villanova University, Villanova, PA 19085, USA*

## Abstract

Let  $v = v(G)$  and  $e = e(G)$  denote the order and size of a simple graph  $G$ , respectively. Let  $\mathcal{G} = \{G_i\}_{i \geq 1}$  be a family of simple graphs of magnitude  $r > 1$  and constant  $\lambda > 0$ , i.e.  $e(G_i) = (\lambda + o(1))v(G_i)^r$ ,  $i \rightarrow \infty$ . For any such family  $\mathcal{G}$  whose members are bipartite and of girth at least  $2k + 2$ , and every integer  $t$ ,  $2 \leq t \leq k - 1$ , we construct a family  $\tilde{\mathcal{G}}_t$  of graphs of same magnitude  $r$ , of constant greater than  $\lambda$ , and all of whose members contain each of the cycles  $C_4, C_6, \dots, C_{2t}$ , but none of the cycles  $C_{2t+2}, \dots, C_{2k}$ . We also prove that for every family of  $2k$ -cycle free extremal graphs (i.e. graphs having the greatest size among all  $2k$ -cycle free graphs of the same order), all but finitely many such graphs must be either non-bipartite or have girth at most  $2k - 2$ . In particular, we show that the best known lower bound on the size of  $2k$ -cycle free extremal graphs for  $k = 3, 5$ , namely  $(2^{-\frac{k+1}{k}} + o(1))v^{\frac{k+1}{k}}$ , can be improved to  $((k - 1) \cdot k^{-\frac{k+1}{k}} + o(1))v^{\frac{k+1}{k}}$ .

## 1. Introduction

All graphs we consider are simple. Let  $\mathcal{F}$  be a family of graphs, and let  $G$  be a graph which contains no subgraph isomorphic to a graph from  $\mathcal{F}$ . Then  $G$  is called  $\mathcal{F}$ -free. We consider the following problem from extremal graph theory: Find the greatest number of edges  $\text{ex}(v, \mathcal{F})$  of any graph on  $v$  vertices which is  $\mathcal{F}$ -free. In this context  $\mathcal{F}$  is called the *family of forbidden graphs*.

Let  $k$  be a fixed but arbitrary integer,  $k \geq 2$ , and let  $C_{2k}$  denote the  $2k$ -cycle. By the Even Circuit Theorem (see [2,4,6]) we have an upper bound for  $\text{ex}(v, \{C_{2k}\})$ :

$$\text{ex}(v, \{C_{2k}\}) = O(v^{1+\frac{1}{k}}).$$

---

<sup>1</sup>This research was supported by NSF grant DMS-9115473.

A general lower bound for  $\text{ex}(v, \{C_{2k}\})$  is also available. In fact [11,12],

$$\text{ex}(v, \{C_{2k}\}) \geq \text{ex}(v, \{C_3, C_4, \dots, C_{2k+1}\}) = \Omega(v^{1+\frac{2}{3k+3}}).$$

For  $3 \leq k \leq 8$ , this bound can be improved [10] to

$$\text{ex}(v, \{C_{2k}\}) \geq \text{ex}(v, \{C_3, C_4, \dots, C_{2k+1}\}) = \Omega(v^{1+\frac{1}{2k-3}}),$$

and for certain values of  $k$  the bound can be improved still further (see [14]). Most notably,  $\text{ex}(v, \{C_4\}) \sim \frac{1}{2}v^{3/2}$  [3,5,7,8] and  $\text{ex}(v, \{C_{2k}\}) = \Omega(v^{1+\frac{1}{k}})$  for  $k = 3$  and  $5$  [1,9,13], but similar results have not been proved for any other values of  $k$ .

Let  $\mathcal{F}$  be a family of graphs, and let  $G$  be an  $\mathcal{F}$ -free graph of order  $v = v(G)$  and size  $e = e(G)$  for which  $e = \text{ex}(v, \mathcal{F})$ . We call such graphs  $G$  *extremal* (or  $\mathcal{F}$ -*extremal* when we wish to emphasize  $\mathcal{F}$ ). Let  $\mathcal{H} = \{H_i\}_{i \geq 1}$  be an arbitrary family of  $\mathcal{F}$ -free graphs such that  $\{v_i = v(H_i)\}_{i \geq 1}$  is an increasing sequence of integers. We denote by  $r(\mathcal{H})$  the least number  $r$  such that  $e(H_i) = O(v_i^r)$ , should such an  $r$  exist, and we call it the *magnitude* of  $\mathcal{H}$ . For a family  $\mathcal{H}$  of magnitude  $r$ , let  $\lambda(\mathcal{H})$  denote the greatest  $\lambda$  such that  $e(H_i) = (\lambda + o(1))v_i^r, i \rightarrow \infty$ , should such a  $\lambda$  exist, and call it the *constant* of  $\mathcal{H}$ . Let  $\mathcal{G} = \{G_i\}_{i \geq 1}$  be any family of  $\mathcal{F}$ -extremal graphs with magnitude  $r(\mathcal{G})$  and constant  $\lambda(\mathcal{G})$ , and  $v_i = v(G_i), i \geq 1$ . Then  $\text{ex}(v_i, \mathcal{F}) = (\lambda(\mathcal{G}) + o(1))v_i^{r(\mathcal{G})}$ . Note that for fixed  $\mathcal{F}$ , all families of extremal graphs have the same magnitude and constant, so we may denote them by  $r_{\mathcal{F}}$  and  $\lambda_{\mathcal{F}}$ . Let us call  $\mathcal{H}$  *magnitude extremal* (or  $\mathcal{F}$ -*magnitude extremal*) if  $r(\mathcal{H}) = r_{\mathcal{F}}$ . In practice, investigation of the constant is limited to magnitude extremal graphs.

As an example, let  $\mathcal{F} = \{C_4\}$ . A family  $\mathcal{H}$  of magnitude extremal graphs consists of bipartite point-line incidence graphs of projective planes of order  $q$  (thick generalized triangles). For this family we have  $r(\mathcal{H}) = 3/2$  and  $\lambda(\mathcal{H}) = 2^{-3/2}$ . But since  $\lambda_{\mathcal{F}} = 1/2$  (see any of [3,5,7,8]) the graphs of  $\mathcal{H}$  are not extremal.

For the remainder of this note we restrict ourselves to the case  $\mathcal{F} = \{C_{2k}\}$ . We also write  $r_k$  and  $\lambda_k$ , should they exist, in place of the more cumbersome  $r_{\{C_{2k}\}}$  and  $\lambda_{\{C_{2k}\}}$ . Note that in this case  $1 < r_k \leq 1 + \frac{1}{k}$ .

We now mention an interesting phenomenon concerning certain families of  $\{C_{2k}\}$ -magnitude extremal graphs. Namely, in every case in which  $r_k$  is known (viz.  $r_2 = 3/2, r_3 = 4/3, r_5 = 6/5$ ) there exists a family of  $\{C_{2k}\}$ -magnitude extremal graphs whose members are bipartite graphs of high girth  $g_k$  (viz.  $g_2 = 6, g_3 = 8, g_5 = 12$ ) [1,9,13]. We think this is quite remarkable since the only requirement of such graphs is that they do not contain  $2k$ -cycles, and it would seem (if not for the aforementioned examples which illustrate otherwise) that magnitude could always be increased by selectively adding edges to form smaller cycles. In this note we show that while  $\{C_{2k}\}$ -magnitude extremality can be achieved with families of bipartite graphs of high girth, ordinary extremality cannot! In particular, we give a general and simple constructive procedure for producing from any family of high girth bipartite  $\{C_{2k}\}$ -magnitude

extremal graphs a family of  $\{C_{2k}\}$ -magnitude extremal graphs with greater constant. More precisely, we prove the following

**Theorem.** *Let  $k \geq 3$  and let  $\mathcal{G}$  be a family of  $2k$ -cycle free graphs with magnitude  $r > 1$  and constant  $\lambda > 0$ , the members of which are bipartite graphs of girth at least  $2k+2$ . Then, for any  $t$ ,  $2 \leq t \leq k-1$ , there exists a family  $\tilde{\mathcal{G}}_t$  of  $2k$ -cycle free graphs with magnitude  $r$  and constant  $\tilde{\lambda} \geq t(\frac{2}{t+1})^r \lambda > \lambda$ , all of whose members are bipartite and contain each of the cycles  $C_4, C_6, \dots, C_{2t}$ , and none of the cycles  $C_{2t+2}, \dots, C_{2k}$ . Consequently, any family of  $\{C_{2k}\}$ -extremal graphs must consist (with finitely many exceptions) either of graphs that are non-bipartite or have girth at most  $2k-2$ .*

## 2. The Construction

Let  $G$  be an arbitrary bipartite graph, the partitions of which we denote by  $P$  (points) and  $L$  (lines) for convenience. For any fixed integer  $t \geq 2$ , construct the bipartite graph  $\tilde{G} = \tilde{G}(t)$  as follows: Let  $P^1, P^2, \dots, P^t$  denote  $t$  disjoint copies of the (labelled) points of  $P$ , i.e.  $P^i = \{p^i | p \in P\}$ ,  $i = 1, 2, \dots, t$ . The vertex and edge sets are now defined by  $V(\tilde{G}) = L \cup P^1 \cup P^2 \cup \dots \cup P^t$  and  $E(\tilde{G}) = \{\{p^i, l\} | \{p, l\} \in E(G), i = 1, 2, \dots, t\}$ .

We will need the following two lemmas.

**Lemma 1.** *Let  $\Delta > 1$  be the maximum degree among all points of bipartite graph  $G$ . Then  $\tilde{G}$  contains each of  $C_4, C_6, \dots, C_{2m}$  as subgraphs, where  $m = \min\{t, \Delta\}$ .*

**Proof.** Let  $p \in P$  have degree  $\Delta$  and let the neighbors of  $p$  be  $l_1, l_2, \dots, l_\Delta \in L$ . Then  $l_1 p^1 l_2 p^2 l_3 p^3 \dots l_i p^i l_1$  is clearly a  $2i$ -cycle,  $2 \leq i \leq m$ .

Let  $\{G_i\}_{i \geq 1}$  be a family of bipartite graphs with magnitude  $r > 1$  and constant  $\lambda > 0$ . Without loss of generality we may assume  $|L| \geq |P|$  for each  $G_i$ . Then, for any  $t \geq 2$ , family  $\{\tilde{G}_i\}_{i \geq 1}$  has magnitude  $\tilde{r} = r$  but the constant  $\tilde{\lambda}$  need not exist. In fact, the existence of  $\tilde{\lambda}$  depends on the behavior of the sequence  $\{\mu_i\}$ , where  $\mu_i$  is defined to be the ratio of the number  $|P|$  of points in  $G_i$  to the number  $v = v(G_i)$  of vertices. Note that  $0 < \mu_i \leq 1/2$ . If  $\mu_i \rightarrow \mu, i \rightarrow \infty$ , then, as we shall show in Lemma 2,  $\tilde{\lambda}$  exists, in fact  $\tilde{\lambda} = t[1 + (t-1)\mu]^{-r} \lambda$ . In any case, we can always find a subsequence of  $\{\mu_i\}$  which converges to some  $\mu, 0 \leq \mu \leq 1/2$ , and hence a subfamily  $\{\tilde{G}_{i(\mu)}\}$  of  $\{\tilde{G}_i\}_{i \geq 1}$  for which  $\tilde{\lambda}$  exists and is described as above. Therefore, without loss of generality, we may assume that  $\mu_i \rightarrow \mu$  for the original family  $\{G_i\}$ .

**Lemma 2.**  *$\{\tilde{G}_i\}$  has magnitude  $\tilde{r} = r$  and constant  $\tilde{\lambda} = t[1 + (t-1)\mu]^{-r} \lambda \geq t(\frac{2}{t+1})^r \lambda$ . Moreover  $\tilde{\lambda} > \lambda$  if  $2 \leq t \leq (r-1)^{-1}$ .*

**Proof.** Clearly, graph  $\tilde{G}_i$  has parameters  $\tilde{v} = v + (t-1)|P|$  and  $\tilde{e} = te$ . Then  $\tilde{e}\tilde{v}^{-r} = te[v + (t-1)|P|]^{-r} = tev^{-r}[1 + (t-1)\mu_i]^{-r} \rightarrow t[1 + (t-1)\mu]^{-r} \lambda := \tilde{\lambda}$ . Since  $1 + (t-1)\mu \leq \frac{1}{2}(t+1)$ , then  $\tilde{\lambda} \geq t(\frac{2}{t+1})^r \lambda$ . The last statement of the lemma follows from the fact that the function  $f(t) = t(\frac{2}{t+1})^r$  is increasing on the interval  $[1, (r-1)^{-1}]$ .

We now proceed to

**Proof of the theorem.** Let  $k \geq 3$  and let  $\mathcal{G}$  be a family of graphs as in the theorem statement. Thus  $G \in \mathcal{G}$  is bipartite and has girth at least  $2k+2$ . For fixed  $t$ ,  $2 \leq t \leq k-1$ , form the family  $\tilde{\mathcal{G}}_t = \{\tilde{G} \mid \Delta(G) \geq t\}$ . (Note that  $\Delta \geq \frac{2e}{v} \sim 2\lambda v^{r-1} \rightarrow \infty$  as  $v \rightarrow \infty$ , so that  $\Delta \geq t$  for  $v$  sufficiently large.) By Lemma 1,  $\tilde{G}$  contains each of the cycles  $C_4, C_6, \dots, C_{2t}$ . Suppose  $\tilde{G}$  contains a  $2s$ -cycle for some  $s$ ,  $t+1 \leq s \leq k$ , which we describe by its sequence of consecutive vertices

$$a_1 b_1 a_2 b_2 \cdots a_s b_s, \quad (1)$$

where  $a_i \in P^1 \cup \dots \cup P^t$ ,  $b_i \in L$ ,  $1 \leq i \leq s$ . We consider the closed walk in  $G$ ,

$$\eta(a_1) b_1 \eta(a_2) b_2 \cdots \eta(a_s) b_s, \quad (2)$$

which is the image of (1) under the map  $\eta : V(\tilde{G}) \rightarrow V(G)$  defined by  $\eta(p^i) = p$  for all  $p^i \in P^i$ ,  $1 \leq i \leq t$ , and  $\eta(l) = l$  for all  $l \in L$ . Note that while the  $b_i$  are certainly distinct, this need not be the case for the  $\eta(a_i)$ .

Sequence (2) defines an eulerian multigraph in which every edge has multiplicity at most 2 and each line  $b_i$  has degree exactly 2. Delete from this multigraph all edges of multiplicity 2 (all 2-cycles). Then, in the resulting simple graph, each connected component is eulerian. But if there exists such a component which is not an isolated vertex, then  $G$  contains a cycle of length at most  $2s$ , a contradiction. Thus, every component is an isolated vertex, i.e. every edge in the multigraph has multiplicity 2. This implies that all  $\eta(a_i)$  are equal, so (1) has the form

$$p^1 b_1 p^2 b_2 \dots p^s b_s$$

for some  $p \in P$ , an impossibility as  $\tilde{G}$  has only  $t$  copies of  $P$  and  $t \leq s-1$ . We conclude that  $\tilde{G}$  is free of all cycles  $C_{2t+2}, \dots, C_{2k}$ . Finally, by Lemma 2,  $\tilde{\mathcal{G}}_t$  has the same magnitude as  $\mathcal{G}$  with constant as in the statement of the theorem. It remains only to show that  $\tilde{\lambda} > \lambda$ . But since  $r \leq 1 + \frac{1}{k}$ , this follows immediately from Lemma 2.

**Corollary.**  $\lambda_3 \geq 2/3^{4/3}, \lambda_5 \geq 4/5^{6/5}$ .

**Proof.** Apply our construction to the known families of magnitude extremal graphs [1,9,13] which have magnitudes  $4/3$  and  $6/5$  and constants  $2^{-4/3}$  and  $2^{-6/5}$ , respectively.

### 3. Concluding Remarks

It would seem that the constructive procedure described in Section 2 could be applied to graphs with forbidden family different from  $\{C_{2k}\}$ . While this is probably the case, the situation might be a bit subtle. For example, consider  $\mathcal{F} = \{K_{3,3}\}$  where  $K_{3,3}$  is the complete bipartite graph on  $3+3$  vertices. It is easy to see that if  $G$  is a

$\{K_{3,3}\}$ -free bipartite graph which just happens to have no  $K_{2,3}$  subgraphs, then  $\tilde{G}(2)$  will be  $\{K_{3,3}\}$ -free and with larger constant. This result shows that any family of  $\{K_{3,3}\}$ -extremal graphs must consist either of non-bipartite graphs or graphs which contain a copy of  $K_{2,3}$ . The point is that, as can easily be shown from a result of Brown [3], *every* family of  $\{K_{3,3}\}$ -magnitude extremal graphs must consist of graphs which contain a  $K_{2,3}$  subgraph, so that, in this case, there are no graphs to which our construction applies.

*Note added in proof.* Recently the authors proved that  $\text{ex}(v, \{C_3, C_4, \dots, C_{2k+1}\}) = \Omega(v^{1+2/(3k-3+\epsilon)})$ , where  $\epsilon = 0$  if  $k$  is odd and  $\epsilon = 1$  if  $k$  is even. To our knowledge this is the best asymptotic lower bound for all  $k$ ,  $k \geq 2$ ,  $k \neq 5$ . (The result will appear elsewhere.)

## References

1. C. T. Benson, Minimal regular graphs of girth eight and twelve, *Canad. J. Math* **18** (1966), 1091–1094.
2. J. A. Bondy and M. Simonovits, Cycles of even length in graphs, *J. Combin. Theory Ser. B* **16** (1974), 87–105.
3. W. G. Brown, On graphs that do not contain a Thomsen graph, *Canad. Math. Bull.* **9**, No. 3 (1966), 281–285.
4. P. Erdős, Extremal problems in graph theory, in *Theory of Graphs and Its Applications*, ed. M. Fiedler (Academic Press, New York, 1985).
5. P. Erdős, A. Rényi and V. T. Sós, On a problem of graph theory, *Studia Sci. Math. Hungar.* **1** (1966), 215–235.
6. R. J. Faudree and M. Simonovits, On a class of degenerate extremal graph problems, *Combinatorica* **3** (1) (1983), 83–93.
7. Z. Füredi, Graphs without quadrilaterals, *J. Combin. Theory, Series B* **34** (1983), 187–190.
8. Z. Füredi, Quadrilateral-free graphs with maximum number of edges, preprint.
9. F. Lazebnik and V. A. Ustimenko, New examples of graphs without small cycles and of large size, *Europ. J. Combin.* **14** (1993), 445–460.
10. F. Lazebnik and V. A. Ustimenko, Explicit construction of graphs with an arbitrary large girth and of large size, *Applied Discrete Math.*, to appear.
11. A. Lubotzky, R. Phillips and P. Sarnak, Ramanujan graphs, *Combinatorica* **8** (3) (1988), 261–277.
12. G. A. Margulis, Explicit group–theoretical construction of combinatorial schemes and their application to the design of expanders and concentrators, *Journal of Problems of Information Transmission*, (1988), 39–46 (translation from *Problemy Peredachi Informatsii*, vol. 24, No. 1, 51–60, January–March 1988).
13. R. Wenger, Extremal graphs with no  $C^4$ ,  $C^6$ , or  $C^{10}$ 's, *J. Combin. Theory, Series B* **52**, 113–116 (1991)
14. A. J. Woldar and V. A. Ustimenko, An application of group theory to extremal graph theory, in “Group Theory: Proceedings of the Biennial Ohio-State Denison Conference” (S. Sehgal and R. Solomon, Eds.), World Scientific Publ. Co., Singapore, 1993.