

Tightness of voter model interfaces

Anja Sturm

Department of Mathematical Sciences
University of Delaware
501 Ewing Hall
Newark, DE 19716-2553, USA
e-mail: sturm@math.udel.edu

Jan M. Swart

ÚTIA
Pod vodárenskou věží 4
18208 Praha 8
Czech Republic
e-mail: swart@utia.cas.cz

July 11, 2007

Abstract

Consider a long-range, one-dimensional voter model started with all zeroes on the negative integers and all ones on the positive integers. If the process obtained by identifying states that are translations of each other is positively recurrent, then it is said that the voter model exhibits interface tightness. In 1995, Cox and Durrett proved that one-dimensional voter models exhibit interface tightness if their infection rates have a finite third moment. Recently, Belhaouari, Mountford, and Valle have improved this by showing that a finite second moment suffices. The present paper gives a new short proof of this fact. We also prove interface tightness for a long range swapping voter model, which has a mixture of long range voter model and exclusion process dynamics.

MSC 2000. Primary: 82C22; Secondary: 82C24, 82C41, 60K35.

Keywords. Long range voter model, swapping voter model, interface tightness, exclusion process.

Acknowledgements. Work sponsored by GAČR grants 201/06/1323 and 201/07/0237, and by UDRF grant 06000596.

1 Introduction and main results

Let $X = (X_t)_{t \geq 0}$ be a long-range, one-dimensional ‘swapping’ voter model, i.e. X is a Markov process with state space $\{0, 1\}^{\mathbb{Z}}$ and formal generator $G := G^v + G^s$, where

$$\begin{aligned} G^v f(x) &:= \sum_{ij} q(i-j) 1_{\{x(i) \neq x(j)\}} \{f(x^{\{i\}}) - f(x)\}, \\ G^s f(x) &:= \frac{1}{2} \sum_{ij} p(i-j) 1_{\{x(i) \neq x(j)\}} \{f(x^{\{i,j\}}) - f(x)\}, \end{aligned} \tag{1.1}$$

q and p are functions $\mathbb{Z} \rightarrow [0, \infty)$, and p is symmetric, i.e., $p(i) = p(-i)$ ($i \in \mathbb{Z}$). Here, for any $x \in \{0, 1\}^{\mathbb{Z}}$ and $\Delta \subset \mathbb{Z}$,

$$x^\Delta(i) := \begin{cases} 1 - x(i) & \text{if } i \in \Delta, \\ x(i) & \text{if } i \notin \Delta \end{cases} \tag{1.2}$$

denotes the configuration obtained from x by flipping all spins in Δ . Note that (1.1) says that if $X_t(i) \neq X_t(j)$ for some $i \neq j$, then due to the action of G^v , $X_t(i)$ adopts the type of $X_t(j)$ with *infection rate* $q(i-j)$. This describes a long-range, one-dimensional voter model. In addition, due to the action of G^s , at *swapping rate* $p(i-j)$ the sites i and j swap their types. In order for the process to be well-defined (see [Lig85]), we assume that $\sum_i (q(i) + p(i)) < \infty$. We will also require irreducibility conditions. We call a rate function $r : \mathbb{Z} \rightarrow [0, \infty)$ *irreducible* if each $i \in \mathbb{Z}$ can be written as $i = i_1 + \dots + i_n$ with $n \geq 0$ and $r(i_k) > 0$ for all $1 \leq k \leq n$.

Consider the space

$$S_{\text{int}} := \{x \in \{0, 1\}^{\mathbb{Z}} : \lim_{i \rightarrow -\infty} x(i) = 0, \lim_{i \rightarrow \infty} x(i) = 1\} \quad (1.3)$$

of states describing the interface between two infinite population of zeroes and ones. If the infection and swapping rates have a finite first moment, i.e., $\sum_i |i|(q(i) + p(i)) < \infty$, then $X_0 \in S_{\text{int}}$ implies $X_t \in S_{\text{int}}$ for all $t \geq 0$, a.s. (see [BMV07] for this statement concerning G^v).

Define an equivalence relation on S_{int} by setting $x \sim y$ if x and y are translations of each other and let $\tilde{S}_{\text{int}} := \{\tilde{x} : x \in S_{\text{int}}\}$, with $\tilde{x} := \{y \in S_{\text{int}} : y \sim x\}$, denote the set of equivalence classes. Then $\tilde{X} = (\tilde{X}_t)_{t \geq 0}$ is a continuous-time Markov process with countable state space \tilde{S}_{int} . Let $x_{\text{H}}(i) := 1_{\{i \geq 0\}}$ denote the ‘Heaviside state’. As long as $q(i) > 0$ for some $i \neq 0$, it is not hard to see that \tilde{x}_{H} can be reached from any state in \tilde{S}_{int} , hence \tilde{X} is an irreducible Markov process on the set of all states that can be reached from \tilde{x}_{H} . Following Cox and Durrett [CD95], we say that X exhibits *interface tightness* if \tilde{X} is positively recurrent on this set.

Theorem 4 in [CD95] states that long-range voter models (without swapping) exhibit interface tightness provided that their infection rates q are symmetric, irreducible, and have a finite third moment. Although not carried out there, their proofs also work if q is asymmetric and q_s is irreducible, where $q_s(i) := \frac{1}{2}(q(i) + q(-i))$ denote the *symmetrized infection rates*. Recently, Belhaouari, Mountford, and Valle [BMV07] improved this result by showing that a finite second moment suffices. They showed that this condition is sharp in the sense that interface tightness does not hold if $\sum_i |i|^c q(i) = \infty$ for some $c < 2$. We will give a new, short proof of their sufficiency result. In fact, we will prove more:

Theorem 1 (Interface tightness for long-range swapping voter models)

Assume that $\sum_i |i|^2(q(i) + p(i)) < \infty$, $q(i) > 0$ for some $i \neq 0$, and that $q_s + p$ is irreducible. Then X exhibits interface tightness.

For the symmetric nearest-neighbor case, Theorem 1 has been proved in [BFMP01, Theorem 1.1 (ii) (b)]. They also have results for asymmetric exclusion dynamics (but symmetric voter dynamics). We note that the symmetric nearest-neighbor swapping voter model arises as the dual and interface model of certain systems of parity preserving branching and annihilating random walks, see [SS07, Section 2.1] and references there.

2 Proof of Theorem 1

The main idea of the proof is the same as in [CD95], namely, to look at the number of ‘inversions’:

$$f_{\text{CD}}(x) := |\{(i, j) \in \mathbb{Z}^2, i < j, x(i) > x(j)\}| \quad (x \in S_{\text{int}}). \quad (2.1)$$

In [CD95, Section 4] and [BMV07], this quantity is estimated using duality and (subtle) results about one-dimensional random walk. Our approach will be to insert f_{CD} into the generator G and prove that if interface tightness would not hold, then f_{CD} would decrease unboundedly as time tends to infinity, which yields a contradiction. In a way, our proof is similar to the methods in [BFMP01], which are based on Lyapunov functions. The function f_{CD} , however, is not a Lyapunov function for our process, so our proofs need a probabilistic ingredient as well, which is provided by Proposition 4 below.

We start by calculating Gf_{CD} .

Lemma 2 (Changes in number of inversions) For each $x \in S_{\text{int}}$, one has

$$Gf_{\text{CD}}(x) = \sum_{n=1}^{\infty} (q_s(n) + p(n))n^2 - \sum_{n=1}^{\infty} q_s(n)I_n(x), \quad (2.2)$$

where

$$I_n(x) := |\{i \in \mathbb{Z} : x(i) \neq x(i+n)\}| \quad (n \geq 1). \quad (2.3)$$

We will need the following lemma:

Lemma 3 (Nonnegative expectation) Assume that $\sum_i |i|^2(q(i) + p(i)) < \infty$. Then

$$\mathbb{E}[f_{\text{CD}}(X_0)] + \int_0^t \mathbb{E}[Gf_{\text{CD}}(X_s)] ds \geq 0 \quad (t \geq 0). \quad (2.4)$$

We will also need a result stating that the number of ‘boundaries’ between zeroes and ones, defined in the sense of (2.3), grows over time if interface tightness does not hold.

Proposition 4 (Interface growth) Assume that $\sum_i |i|(q(i) + p(i)) < \infty$, $q(i) > 0$ for some $i \neq 0$, and $q_s + p$ is irreducible. Assume that interface tightness does not hold. Then

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \mathbb{P}[I_n(X_t) < N] = 0 \quad (N, n \geq 1). \quad (2.5)$$

With these statements we are now in the position to prove Theorem 1. Assume that interface tightness does not hold. By our assumptions on q and p we can choose $i, N \geq 1$ such that

$$\sum_{n=1}^{\infty} (q_s(n) + p(n))n^2 < q_s(i)N. \quad (2.6)$$

Then, by Lemmas 3 and 2, and Proposition 4, the process started in $X_0 = x_H$ satisfies

$$\begin{aligned} 0 &\leq \int_0^T dt \mathbb{E}[Gf_{\text{CD}}(X_t)] \\ &= T \sum_{n=1}^{\infty} (q_s(n) + p(n))n^2 - \sum_{n=1}^{\infty} q_s(n) \int_0^T dt \mathbb{E}[I_n(X_t)] \\ &\leq T \sum_{n=1}^{\infty} (q_s(n) + p(n))n^2 - q_s(i)N \int_0^T dt \mathbb{P}[I_i(X_t) \geq N] \\ &= T \sum_{n=1}^{\infty} (q_s(n) + p(n))n^2 - (T - o(T))q_s(i)N, \end{aligned} \quad (2.7)$$

as $T \rightarrow \infty$. Due to (2.6) the right-hand side of (2.7) tends to $-\infty$ as $T \rightarrow \infty$, which yields a contradiction. \blacksquare

3 Proof of Lemmas 2 and 3, and Proposition 4

Proof of Lemma 2 We need to count the number of pairs of sites i, j with $i < j$ and $x(i) > x(j)$ that are created and deleted due to the various possible jumps. We will first consider $G^s f_{CD}$, i.e. the changes due to swapping. So consider the case that $x(i) \neq x(i+n)$ for some $n \in \mathbb{N}$, while there are l ones on the left of i , r zeroes on the right of $i+n$, and n_0 zeroes and n_1 ones between i and $i+n$. Then the changes in $f_{CD}(x)$ due to swapping can be summarized as follows:

$$\begin{aligned} \underbrace{\dots}_{l \times 1} \quad 0 \quad \underbrace{\dots}_{n_0 \times 0, n_1 \times 1} \quad 1 \quad \underbrace{\dots}_{r \times 0} &\rightarrow \dots 1 \dots 0 \dots \quad n_0 + n_1 + 1 = n \\ \underbrace{\dots}_{l \times 1} \quad 1 \quad \underbrace{\dots}_{n_0 \times 0, n_1 \times 1} \quad 0 \quad \underbrace{\dots}_{r \times 0} &\rightarrow \dots 0 \dots 1 \dots \quad -(n_0 + n_1 + 1) = -n \end{aligned} \quad (3.1)$$

Thus, if we define

$$I_n^{ab}(x) := |\{i : x(i) = a, x(i+n) = b\}| \quad (n \geq 0, ab = 01, 10) \quad (3.2)$$

then we obtain

$$G^s f_{CD}(x) = \sum_{n=1}^{\infty} p(n)(n \cdot I_n^{01}(x) - n \cdot I_n^{10}(x)) \quad (3.3)$$

Now, for any $0 \leq m < n$, set

$$I_{n,m}^{ab}(x) := |\{r \in \mathbb{Z} : x(nr+m) = a, x(n(r+1)+m) = b\}|. \quad (3.4)$$

Walking along the thinned-out lattice $n\mathbb{Z} + m$ from $-\infty$ to $+\infty$, we see one more change from 0 to 1 than we see changes from 1 to 0, i.e., $I_{n,m}^{01}(x) = I_{n,m}^{10}(x) + 1$ for all $x \in S_{\text{int}}$. Since $I_n^{ab}(x) = \sum_{m=0}^{n-1} I_{n,m}^{ab}(x)$, it follows that

$$I_n^{01}(x) = I_n^{10}(x) + n \quad (x \in S_{\text{int}}). \quad (3.5)$$

This implies that (3.3) simplifies to

$$G^s f_{CD}(x) = \sum_{n=1}^{\infty} p(n)n^2. \quad (3.6)$$

In order to consider the effect of G^v on f_{CD} we write

$$\begin{aligned} G^v f_{CD}(x) &= \sum_k q(-k) \left(\sum_{i: x(i)=1} \sum_{j: j>i} (1_{\{x(j)=1, x(j+k)=0\}} - 1_{\{x(j)=0, x(j+k)=1\}}) \right. \\ &\quad \left. + \sum_{i: x(i)=0} \sum_{j: j<i} (1_{\{x(j)=0, x(j+k)=1\}} - 1_{\{x(j)=1, x(j+k)=0\}}) \right). \end{aligned} \quad (3.7)$$

We observe that for any $x \in S_{\text{int}}$ and $k > 0$,

$$\begin{aligned} &\sum_{j: j>i} (1_{\{x(j)=1, x(j+k)=0\}} - 1_{\{x(j)=0, x(j+k)=1\}}) \\ &= \sum_{j=i+1}^{i+k} \sum_{n \geq 0} (1_{\{x(j+nk)=1, x(j+(n+1)k)=0\}} - 1_{\{x(j+nk)=0, x(j+(n+1)k)=1\}}) \\ &= - \sum_{j=i+1}^{i+k} 1_{\{x(j)=0\}}. \end{aligned} \quad (3.8)$$

To see why the last equality in (3.8) holds, observe that since $k > 0$, the sequence $x(j), x(j+k), x(j+2k) \dots$ is eventually one. Hence, if $x(j) = 1$, then the number of changes from 0 to 1 equals the number of changes from 1 to 0 and all terms cancel, while if $x(j) = 0$ there is one extra change from 0 to 1, leading to a contribution of minus one.

Likewise, for any $x \in S_{\text{int}}$ and $k > 0$,

$$\begin{aligned}
& \sum_{j:j < i} \left(\mathbf{1}_{\{x(j)=0, x(j+k)=1\}} - \mathbf{1}_{\{x(j)=1, x(j+k)=0\}} \right) \\
&= \sum_{j=i-k}^{i-1} \sum_{n \geq 0} \left(\mathbf{1}_{\{x(j-nk)=0, x(j-(n-1)k)=1\}} - \mathbf{1}_{\{x(j-nk)=1, x(j-(n-1)k)=0\}} \right) \\
&= \sum_{j=i}^{i+k-1} \sum_{n \geq 0} \left(\mathbf{1}_{\{x(j-(n+1)k)=0, x(j-nk)=1\}} - \mathbf{1}_{\{x(j-(n+1)k)=1, x(j-nk)=0\}} \right) \\
&= \sum_{j=i}^{i+k-1} \mathbf{1}_{\{x(j)=1\}}.
\end{aligned} \tag{3.9}$$

It follows that for any $k > 0$,

$$\begin{aligned}
& \sum_{i:x(i)=1} \sum_{j:j > i} \left(\mathbf{1}_{\{x(j)=1, x(j+k)=0\}} - \mathbf{1}_{\{x(j)=0, x(j+k)=1\}} \right) \\
&+ \sum_{i:x(i)=0} \sum_{j:j < i} \left(\mathbf{1}_{\{x(j)=0, x(j+k)=1\}} - \mathbf{1}_{\{x(j)=1, x(j+k)=0\}} \right) \\
&= - \sum_i \sum_{j=i+1}^{i+k} \mathbf{1}_{\{x(i)=1, x(j)=0\}} + \sum_i \sum_{j=i}^{i+k-1} \mathbf{1}_{\{x(i)=0, x(j)=1\}} \\
&= \sum_{n=1}^{k-1} I_n^{01}(x) - \sum_{n=1}^k I_n^{10}(x),
\end{aligned} \tag{3.10}$$

Using (3.5), the expression in (3.10) can be rewritten as

$$- I_k^{10}(x) + \sum_{n=1}^{k-1} (I_n^{01}(x) - I_n^{10}(x)) = -\frac{1}{2}(I_k(x) - k) + \sum_{n=1}^{k-1} n = \frac{1}{2}(k^2 - I_k(x)), \tag{3.11}$$

which holds for $k > 0$. Using symmetry with respect to the map $x \mapsto x'$ where $x'(i) := 1 - x(-i)$, it is not hard to see that we get the same formula for the expression in (3.10) if $k < 0$. Inserting this into (3.7), we arrive at

$$G^v f_{\text{CD}}(x) = \sum_{n=1}^{\infty} q_s(n)(n^2 - I_n(x)). \tag{3.12}$$

Taking (3.6) and (3.12) together now implies (2.2). ■

Proof of Lemma 3 If the function f_{CD} were bounded, then standard theory would tell us that the process

$$M_t := f_{\text{CD}}(X_t) - \int_0^t G f_{\text{CD}}(X_s) ds \quad (t \geq 0) \tag{3.13}$$

is a martingale with respect to the filtration generated by X . In particular, since $\mathbb{E}[M_t] = \mathbb{E}[M_0]$ and $f_{\text{CD}} \geq 0$, this would imply (2.4). In the present case, since f_{CD} is unbounded, we have to work a bit. Let

$$w(x) := \max\{i : x(i) \neq x(i+1)\} - \min\{i : x(i) \neq x(i+1)\} \quad (3.14)$$

denote the ‘width’ of an interface state $x \in S_{\text{int}}$. We can couple X_t to a continuous-time random walk $(R_t)_{t \geq 0}$, started in $R_0 = w(X_0)$, which jumps from r to $r+n$ with rate

$$a(n) := \sum_{k=n}^{\infty} 2(q_s(k) + p(k)), \quad (3.15)$$

in such a way that $R_t \geq w(X_t)$ for all $t \geq 0$ a.s. We check that

$$\begin{aligned} \sum_{n=1}^{\infty} a(n)n &= \sum_{n=1}^{\infty} \sum_{k=n}^{\infty} 2(q_s(k) + p(k))n = \sum_{k=1}^{\infty} \sum_{n=1}^k 2(q_s(k) + p(k))n \\ &= \sum_{k=1}^{\infty} 2(q_s(k) + p(k)) \frac{1}{2} k(k+1) < \infty. \end{aligned} \quad (3.16)$$

Let X^K be the process with generator given by

$$\begin{aligned} G^K f(x) &:= \sum_{ij: |i-j| \leq K} q(i-j) 1_{\{x(i) \neq x(j)\}} \{f(x^{\{i\}}) - f(x)\}, \\ &\quad + \frac{1}{2} \sum_{ij: |i-j| \leq K} p(i-j) 1_{\{x(i) \neq x(j)\}} \{f(x^{\{i,j\}}) - f(x)\}, \end{aligned} \quad (3.17)$$

started in $X_0^K = X_0$. We set

$$\tau_{K,N} := \inf\{t \geq 0 : w(X_t^K) > N\} \quad (3.18)$$

and define τ_N similarly, with X^K replaced by X . It follows from standard theory that

$$M_t^{K,N} := f_{\text{CD}}(X_{t \wedge \tau_{K,N}}^K) - \int_0^{t \wedge \tau_{K,N}} G^K f_{\text{CD}}(X_s^K) ds \quad (t \geq 0) \quad (3.19)$$

is a martingale, where by Lemma 2,

$$G^K f_{\text{CD}}(x) = \sum_{n=1}^K (q_s(n) + p(n))n^2 - \sum_{n=1}^K q_s(n)I_n(x). \quad (3.20)$$

Using an obvious coupling, we may arrange that X^K converges a.s. to X in such a way that there exists a random K_0 such that for all $K \geq K_0$, one has $\tau_{K,N} = \tau_N$ and $X_t^K = X_t$ for all $t \leq \tau_N$. Since moreover $\tau_N \rightarrow \infty$ as $N \rightarrow \infty$ (which follows from the estimate $w(X_t) \leq R_t$), taking into account (2.2) and (3.20), it follows that

$$\lim_{N \rightarrow \infty} \lim_{K \rightarrow \infty} \int_0^{t \wedge \tau_{K,N}} G^K f_{\text{CD}}(X_s^K) ds = \int_0^t G f_{\text{CD}}(X_s) ds \quad \text{a.s.} \quad (3.21)$$

Since $M^{K,N}$ is a martingale,

$$0 \leq \mathbb{E}[f_{\text{CD}}(X_{t \wedge \tau_{K,N}}^K)] = \mathbb{E}[f_{\text{CD}}(X_0)] + \mathbb{E}\left[\int_0^{t \wedge \tau_{K,N}} G^K f_{\text{CD}}(X_s^K) ds\right]. \quad (3.22)$$

Letting $K \rightarrow \infty$ and then $N \rightarrow \infty$ in (3.22), using (3.21), we arrive at (2.4), provided we show (in view of (3.20)) that the random variables

$$\sum_{n=1}^K q_s(n) \int_0^{t \wedge \tau_{K,N}} I_n(X_s^K) ds \quad (K, N \geq 1) \quad (3.23)$$

are uniformly integrable. We can couple the X^K to a random walk $(R_t)_{t \geq 0}$ with jump rates $a(n)$ as in (3.15), in such a way that $R_t \geq w(X_t^K)$ for all $t \geq 0$ and $K \geq 1$ a.s. Hence, since $I_n(x) \leq n + w(x)$, we may estimate

$$\sum_{n=1}^K q_s(n) \int_0^{t \wedge \tau_{K,N}} I_n(X_s^K) ds \leq \sum_{n=1}^{\infty} q_s(n) \int_0^t (n + R_s) ds. \quad (3.24)$$

By (3.16), the right-hand side of (3.24) has finite expectation, proving the required uniform integrability. \blacksquare

Remark 5 (Martingale problem) If $\sum_i |i|^3 (q(i) + p(i)) < \infty$, then the jump rates $a(n)$ in (3.15) have a finite second moment. Using this and the estimate $f_{\text{CD}}(x) \leq w(x)^2$, one can prove that in this case the process in (3.13) is a martingale.

Proof of Proposition 4 Consider the ‘boundary process’

$$Y_t(i) := 1_{\{X_t(i) \neq X_t(i+1)\}} \quad (t \geq 0, i \in \mathbb{Z}), \quad (3.25)$$

which is a Markov process in

$$S_{\text{bound}} := \{y \in \{0, 1\}^{\mathbb{Z}} : \sum_i y(i) \text{ is finite and odd}\}. \quad (3.26)$$

Then $(Y_t)_{t \geq 0}$ is a Markov process in $\{0, 1\}^{\mathbb{Z}}$ with formal generator

$$\begin{aligned} G_Y f(y) := & \sum_{i < j} 1_{\{\sum_{k=i+1}^j y(k) \text{ is odd}\}} q(i-j) \{f(y^{\{i, i+1\}}) - f(y)\} \\ & + \sum_{i > j} 1_{\{\sum_{k=j+1}^i y(k) \text{ is odd}\}} q(i-j) \{f(y^{\{i, i+1\}}) - f(y)\} \\ & + \sum_{i < j-1} 1_{\{\sum_{k=i+1}^j y(k) \text{ is odd}\}} p(i-j) \{f(y^{\{i, i+1, j, j+1\}}) - f(y)\} \\ & + \sum_i 1_{\{y(i+1) = 1\}} p(1) \{f(y^{\{i, i+2\}}) - f(y)\}. \end{aligned} \quad (3.27)$$

Using the fact that $\sum_i |i|(q(i) + p(i)) < \infty$, one can check that Y is a parity preserving cancellative spin system in the sense of [SS07]. We will show that the process started in y , denoted by Y^y , satisfies

$$\inf \{ \mathbb{P}[|Y_t^y| = n] : |y| = n + 2, y(i) = 1 = y(j) \text{ for some } i \neq j, |i - j| \leq L \} > 0 \quad (3.28)$$

($L \geq 1, n \geq 0, t > 0$). As a result of (3.28), we can apply [SS07, Proposition 13] to conclude that

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \mathbb{P}[I_1(X_t) < N] = 0 \quad (N \geq 1). \quad (3.29)$$

See also Proposition 2.6 in [Han99] for similar arguments concerning a dual process to the threshold voter model.

To give a rough idea of the proof and of the importance of condition (3.28), think of the sites with $Y_t(i) = 1$ as being occupied by a particle. Then Y is a parity preserving particle system, i.e., if Y is started in an odd (even) initial state, then the number of particles always stays odd (even). Let \tilde{Y} denote the process obtained from Y by identifying states that are a translation of each other. If \tilde{X} is not positively recurrent, then the same is true for \tilde{Y} . The main idea of the proof of (3.29) is to use induction on n to show that there cannot be less than n particles for a positive fraction of time. This is obviously true for $n = 1$; imagine that it holds for a certain n . If we see n particles for a positive fraction of time, then most of the time these particles must be situated far from each other, for else with positive probability at least two would annihilate each other due to (3.28), violating the induction hypothesis. However, since \tilde{Y} is not positively recurrent, n single particles far from each other will soon each produce three particles, hence we cannot see n particles for a positive fraction of time.

To boost up (3.29) to the statement in (2.5), it suffices to show that the process started in x , denoted by X^x , satisfies

$$\lim_{N \rightarrow \infty} \inf_{|I_1(x)| \geq N} \mathbb{P}[I_n(X_t^x) < M] = 0 \quad (M, n \geq 1, t > 0). \quad (3.30)$$

For if (3.30) holds, then for each $t, \varepsilon > 0$ we can choose N large enough such that the limit in (3.30) is smaller than ε , and therefore, by (3.29) and a restart argument

$$\limsup_{T \rightarrow \infty} \frac{1}{T} \int_t^{t+T} ds \mathbb{P}[I_n(X_s) < M] \leq \varepsilon \quad (M \geq 1). \quad (3.31)$$

Since $\varepsilon > 0$ is arbitrary, this implies (2.5).

We still need to prove (3.28) and (3.30). We start with the former. Choose $k \neq 0$ such that $q(k) > 0$. By symmetry, we may without loss of generality assume that $k > 0$. It suffices to prove (3.28) for $L \geq k$. So fix $L \geq k$, $n \geq 0$, and $t > 0$. Let $y = \delta_{i_1} + \dots + \delta_{i_n}$ where $\delta_i(j) := 1$ if $i = j$ and $= 0$ otherwise, and $i_1 < \dots < i_n$. Assume that

$$M := \{m' \in \{1, \dots, n-1\} : i_{m'+1} - i_{m'} \leq L\} \quad (3.32)$$

is not empty and let $m := \inf(M)$. Then $x(i_m + 1 - L) = \dots = x(i_m) \neq x(i_m + 1)$, and hence, since $L \geq k$, there is a positive probability that during the time interval $[0, t]$, the sites $i_m + 1, \dots, i_{m+1}$ get infected successively by the sites $i_m + 1 - k, \dots, i_{m+1} - k$, leading to a decrease in $I_1(X_t)$ of 2. Using the fact that $\sum_i |i|(q(i) + p(i)) < \infty$, it is not hard to see that moreover, with positive probability, no other infections take place, and that this probability is uniformly bounded from below in all y satisfying our assumptions.

To prove (3.30), we view the dynamics of our process X as follows. For each ordered pair (i, j) with $i \neq j$, at times selected according to an independent Poisson point process with intensity $q(i - j)$, the type of site j infects the site i . Likewise, for each unordered pair $\{i, j\}$ with $i \neq j$, at times selected according to an independent Poisson point process with intensity $p(i - j)$, the sites i and j swap their types.

We claim that if we view the evolution of types in this way, then we can find a $j \in \mathbb{Z}$ such that with positive probability $X_t(j)$ inherits its type from $x(0)$ and $X_t(j + n)$ inherits its type from $x(1)$. To see this, we will make a number of infections and swaps to transport the type of site 0 to j and the type of site 1 to $j + n$. Let us say that in the k -th step of our construction,

we have transported the type of site 0 to the site l_k and the type of site 1 to r_k . Then, in the $(k + 1)$ -th step of our construction, by making an infection with rate $q(i)$, we may transport the type of site l_k to $l_k + i$ and set $(l_{k+1}, r_{k+1}) := (l_k + i, r_k)$, or we may transport the type of site r_k to $r_k + i$ and set $(l_{k+1}, r_{k+1}) := (l_k, r_k + i)$, and similarly for swaps. Thus, in each step, we may increase $r_k - l_k$ by d for each $d \in \mathcal{G} := \{i \in \mathbb{Z} : q_s(i) + p(i) > 0\}$. We must only make sure that when we move the type of one site to a new position, we do not influence the type of the other site. To avoid this sort of influence, we will make sure that at each point in our construction $l_k < r_k$. We claim that this is possible. Set $n_k := r_k - l_k$. We need to show that there exist (*strictly*) positive integers n_0, \dots, n_m such that $n_0 = 1$, $n_m = n$, and $(n_k - n_{k-1}) \in \mathcal{G}$ for all $1 \leq k \leq m$. By our assumption that $q_s + p$ is irreducible we can write $n - 1 = i_1 + \dots + i_m$ with $i_1, \dots, i_m \in \mathcal{G}$. (Note that this is the only place in our proofs where we use irreducibility.) Without loss of generality we may assume that $i_1 \geq \dots \geq i_m$. Then setting $n_k := 1 + i_1 + \dots + i_k$ proves our claim.

It follows that there exist $j \in \mathbb{Z}$ and $L \geq j$ such that whenever $x(i) \neq x(i + 1)$, there is a positive probability that $X_t(i + j)$ inherits its type from $x(i)$ and $X_t(i + j + n)$ inherits its type from $x(1)$ through a sequence of infections and swaps that are entirely contained in $\{i - L, \dots, i + L\}$. If $I_1(x)$ is large, we can find many sites i , situated at least a distance $3L$ from each other, such that $x(i) \neq x(i + 1)$. By what we have just proved each pair has an independent probability to produce at time t two sites $i + j$ and $i + j + n$ such that $X_t(i + j) \neq X_t(i + j + n)$, hence $I_n(X_t)$ is with large probability large.

Acknowledgement We thank Rongfeng Sun who found an error in our original statement of Lemma 3 and told us how to fix it.

References

- [BFMP01] V. Belitsky, P.A. Ferrari, M.V. Menshikov, and S.Y. Popov. A mixture of the exclusion process and the voter model. *Bernoulli* 7(1): 119–144, 2001.
- [BMV07] S. Belhaouari, T. Mountford, and G. Valle. Tightness for the interfaces of one-dimensional voter models. *Proc. London Math. Soc.* 94(3): 421–442, 2007.
- [CD95] J.T. Cox and R. Durrett. Hybrid zones and voter model interfaces. *Bernoulli* 1(4): 343–370, 1995.
- [Han99] S.J. Handjani. The complete convergence theorem for coexisting threshold voter models. *Ann. Probab.* 27(1): 226–245.
- [Lig85] T.M. Liggett. *Interacting Particle Systems*. Springer, New York, 1985.
- [SS07] A. Sturm and J.M. Swart. Voter models with heterozygosity selection. Accepted in *Ann. Appl. Probab.* ArXiv: math.PR/0701555v2.